

On the quantum super Virasoro algebra

M. Mansour

Laboratoire de Physique Théorique, Faculté des Sciences
B. P. 1014 Rabat. Morocco *

February 7, 2008

ABSTRACT

The quantum super-algebra structure on the deformed super Virasoro algebra is investigated. More specifically we established the possibility of defining a non trivial Hopf super-algebra on both one and two-parameters deformed super Virasoro algebras.

*e-mail: mansour70@mailcity.com

Quantum algebras (quantum groups)[1, 2], are generalizations of the usual Lie algebras to which they reduce for appropriate values of the deformation parameters. From the mathematical point of view they are Hopf algebras [3].

Quantum superalgebras appeared naturally when the quantum inverse scattering method [4] was generalized to the super-systems [4]. Related R-matrix were considered in [5, 6] and simple examples were presented in [7]. The works [8, 9, 10] are devoted to the q-bosonization of the q-superalgebras. The simple example of the dual objects, i.e. quantum supergroups, were investigated in [10, 11, 12]. A complete classification of finite dimensional Lie Superalgebras over C has been given by Kac [13, 14] and Scheunert [15]. Supersymmetry seems to be in the realm of almost all attempts to obtain a unified theory of all interactions [17, 18]. In particular, many problems of string theory are solved once supersymmetry is introduced [19, 20]. The interplay of supersymmetry and conformal invariance is an interesting application of the rules up to now. The two-dimensional superconformal field theory can be treated from a group theoretical point of view, the basic ingredient is the two-dimensional superconformal algebra, which is infinite dimensional called the super Virasoro, see [21] and references therein. The central extension of the super Virasoro algebra leads to the so called the Ramond-Neveu-Schwarz algebras[22, 23]. A supersymmetric extension of the work of Curtright and Zachos is given in [24] by using a quantum superspace approach [25, 26].

The use of quantum superalgebras in physics became popular with the introduction of the deformed harmonic oscillator as a tool for providing a realization of the quantum (super) algebra [27, 28, 7]. Particularly, a great deal of attention has been paid to the deformation of the Virasoro algebra [29, 30, 31, 32, 33, 34] and the super Virasoro algebra [35, 36, 24]. Several investigations of the latter super-algebra are based on the coordinates and derivatives on super space [24, 35]. The quantum group structure on the Deformed Virasoro algebra is given in [37, 38]. However, the existence of a quantum super-algebra structure of the deformed super Virasoro algebra is still an open question. The purpose of this paper is to give a realization of the deformed super Virasoro algebra in terms of bosonic and fermionic algebra operators and define a quantum super-algebra structure on this deformed super-algebra. In ref. [39], an algebraic deformation (associated to an R-matrix of unit square) which allows to a non trivial Hopf super-algebra structure on the super Virasoro and Ramond-Neveu-Schwarz algebras will be

investigated.

First, let us recall that the classical super Virasoro algebra ($sVir$) is defined as an infinite super-algebra generated by the generators $\{L_n, G_n, F_n; n \in \mathbb{Z}\}$ satisfying the defining relations [19, 21]

$$\begin{aligned} [L_n, L_m] &= (n - m)L_{n+m}, \\ [F_m, G_n] &= G_{n+m}, \\ [L_n, F_m] &= -mF_{n+m}, \\ [F_m, F_n] &= 0, \\ [L_m, G_n] &= (m - n)G_{n+m}, \\ [G_m, G_n] &= 0, \end{aligned} \tag{1}$$

and that the Hopf super-algebra structure on the enveloping super-algebra of this classical super-algebra $U(sVir)$ is trivial. The $\mathbb{Z}_2$ -grading on this super-algebra is defined by requiring that $\deg(L_i) = \deg(F_i) = 0$ and $\deg(G_i) = 1$ for $(i \in \mathbb{Z})$. Further, the bracket $[\cdot, \cdot]$ in (1) stands for a graded one

$$[x, y] = xy - (-1)^{\deg(x)\deg(y)}yx.$$

A possible realization of the classical super Virasoro algebra (1) is given as follows

$$\begin{aligned} L_k &= -(a_+)^{k+1}a_-, \\ G_n &= (a_+)^{n+1}f_+a_-, \\ F_n &= (a_+)^nf_+f_- \end{aligned} \tag{2}$$

where the operators $\{a_+, a_-, 1\}$ generate the classical bosonic algebra

$$\begin{aligned} a_-a_+ - a_+a_- &= 1 \\ [1, a_{\pm}] &= 0. \end{aligned} \tag{3}$$

and the remaining ones $\{f_+, f_-, 1\}$ generate the classical fermionic algebra given by

$$\begin{aligned} f_-f_+ + f_+f_- &= 1 \\ [1, f_{\pm}] &= 0, \quad (f_{\pm})^2 = 0. \end{aligned} \tag{4}$$

A q-deformed version of the super Virasoro algebra will be defined by the

generators $\{L_n, G_n, K_n; n \in \mathbb{Z}\}$ and the following q -deformed relations

$$\begin{aligned}
q^{l-k}L_lL_k - q^{k-l}L_kL_l &= [l-k]L_{k+l} \\
F_mG_n - G_nF_m &= G_{n+m}, \\
L_lF_k - q^{2k}F_kL_l &= -q[[k]]F_{k+l}, \\
q^{n-m}F_mF_n - q^{m-n}F_nF_m &= \lambda[n-m]F_{n+m}, \\
q^{l-k}L_lG_k - q^{k-l}G_kL_l &= [l-k]G_{k+l}, \\
G_mG_n + G_nG_m &= 0
\end{aligned} \tag{5}$$

(where $[x] = \frac{q^x - q^{-x}}{q - q^{-1}}$, $[[x]] = \frac{1 - q^{2x}}{1 - q^2}$ and $\lambda = q - q^{-1}$).

The q -deformed super-algebra (5) can be realized in terms of a q -deformed bosonic algebra generators $\{a_+, a_-, 1\}$ and a classical fermionic algebra generators $\{f_+, f_-, 1\}$ which are such that

$$\begin{aligned}
[a_-, a_+]_{q^2} &= 1, & \{f_-, f_+\} &= 1 \\
[a_{\pm}, f_{\pm}] &= 0, & [a_{\mp}, f_{\pm}] &= 0, \\
(f_-)^2 &= 0, & (f_+)^2 &= 0, \\
[1, a_{\pm}] &= 0, & [1, f_{\pm}] &= 0
\end{aligned} \tag{6}$$

(where $[a, b]_{\alpha} = ab - \alpha ba$).

In fact, the following operators

$$\begin{aligned}
L_k &= -q(a_+)^{k+1}a_-, \\
G_n &= (a_+)^{n+1}f_+a_-, \\
F_n &= (a_+)^nf_+f_-
\end{aligned} \tag{7}$$

generate the q -deformed super Virasoro algebra (5).

Now we define the quantum super-algebra structure on the q -deformed enveloping super-algebra of the q -deformed super Virasoro algebra $U_q(sVir_q)$ as follows:

$$\begin{aligned}
\Delta(L_i) &= L_i \otimes T_i + T_i \otimes L_i, \\
\Delta(G_i) &= G_i \otimes R_i + R_i \otimes G_i, \\
\Delta(F_i) &= F_i \otimes K_i + K_i \otimes F_i, \\
\epsilon(L_i) &= 0, \\
\epsilon(G_i) &= 0, \\
\epsilon(F_i) &= 0, \\
S(L_i) &= -S(T_i)L_iT_i^{-1}, \\
S(G_i) &= -S(R_i)G_iR_i^{-1}, \\
S(F_i) &= -S(K_i)F_iK_i^{-1},
\end{aligned} \tag{8}$$

where the operators $\{K_i, T_i, R_i, \quad i \in Z\}$ are additional invertible even elements of $U_q(sVir_q)$ which reduce to the unity operator when the deformation parameter $q = 1$.

From the consistence of the costructures on q -deformed super Virasoro algebra generators with the basic axioms of the Hopf super-algebra structure (co-associativity, co-unity and antipode), we deduce that the costructures on the additional elements are given by

$$\begin{aligned}
\Delta(T_i) &= T_i \otimes T_i, \\
\Delta(R_i) &= R_i \otimes R_i, \\
\Delta(K_i) &= K_i \otimes K_i, \\
\epsilon(T_i) &= 1, \\
\epsilon(R_i) &= 1, \\
\epsilon(K_i) &= 1, \\
S(T_i) &= T_i^{-1}, \\
S(R_i) &= R_i^{-1}, \\
S(K_i) &= K_i^{-1}.
\end{aligned} \tag{9}$$

Further, the consistence of the Hopf super-algebra structure (8) with the deformed commutations relations (5), (i.e. using the fact that the coproduct (Δ) and counit (ϵ) operations must be super-algebra homomorphisms, and that the coinverse (S) must be super-algebra anti-homomorphism) we obtain the following relations between the q -deformed super Virasoro algebra generators and the additional elements.

$$\begin{aligned}
L_k T_l &= q^{2(k+1)l} T_l L_k, \\
L_l K_k &= q^{-1k} K_k L_l, \\
G_k K_l &= K_l G_k, \\
T_l F_k &= q^{(2+1)k} F_k T_l, \\
F_k R_l &= R_l F_k, \\
G_k T_l &= q^{2(1+k)l} T_l G_k, \\
L_l R_k &= q^{2(l+l)k} R_k L_l, \\
F_k K_l &= q^{-2(k+1)l} K_l F_k, \\
G_k R_l &= R_l G_k,
\end{aligned} \tag{10}$$

where the additional generators must satisfy the following relations between

them

$$\begin{aligned}
T_k T_l &= T_l T_k = T_{k+l}, \\
T_k K_l &= K_l T_k = K_{l+k} \\
R_k K_l &= K_l R_k, \\
K_k K_l &= K_l K_k = K_{l+k}, \\
R_k R_l &= R_l R_k = R_{k+l}, \\
T_l R_k &= R_k T_l = R_{k+l}.
\end{aligned} \tag{11}$$

Considering now the two-parameters (p, q) -deformation version of the super Virasoro algebra given by

$$\begin{aligned}
q^{l-k} L_l L_k - q^{k-l} L_k L_l &= [l-k] L_{k+l}, \\
F_m G_n - G_n F_m &= G_{n+m}, \\
L_l F_k - q^{2k} F_k L_l &= -q[[k]] F_{k+l}, \\
q^{n-m} F_m F_n - q^{m-n} F_n F_m &= \lambda[n-m] F_{n+m}, \\
q^{l-k} L_l G_k - p^{2l} q^{k-l} G_k L_l &= [l-k] G_{k+l}, \\
G_m G_n + G_n G_m &= 0
\end{aligned} \tag{12}$$

which can be realized as

$$\begin{aligned}
L_k &= -q(a_+)^{k+1} a_-, \\
G_n &= p^{-2} (a_+)^{n+1} f_+ a_-, \\
F_n &= (a_+)^n f_+ f_-,
\end{aligned} \tag{13}$$

where the generators $\{a_+, a_-, f_+, f_-, 1\}$ satisfy now

$$\begin{aligned}
[a_-, a_+]_{q^2} &= 1, \quad \{f_-, f_+\} = 1 \\
[a_+, f_+]_{p^2} &= 0, \quad [a_-, f_+]_{p^{-2}} = 0, \\
[a_-, f_-]_{p^2} &= 0, \quad [a_+, f_-]_{p^{-2}} = 0, \\
(f_-)^2 &= 0, \quad (f_+)^2 = 0. \\
[1, a_{\pm}] &= 0, \quad [1, f_{\pm}] = 0.
\end{aligned} \tag{14}$$

In this case, the (p, q) -quantum super-algebra structure on the $U_{p,q}(sVir_{p,q})$, will be given by the same costructures on the q -deformed super Virasoro algebra generators (8) and with the same costructures on the additional even generators. But the deformed relations between the (p, q) -deformed super

Virasoro algebra generators and the additional elements are

$$\begin{aligned}
L_k T_l &= q^{2(k+1)l} T_l L_k, \\
L_l K_k &= q^{(1+l)k} K_k L_l, \\
G_k K_l &= K_l G_k, \\
F_k T_l &= q^{(k-1)l} T_l F_k, \\
F_k R_l &= R_l F_k, \\
T_l G_k &= q^{-2(1+k)l} p^{(k+2)l} G_k T_l, \\
L_l R_k &= q^{2(l+l)k} p^{-kl} R_k L_l, \\
F_k K_l &= q^{-2(k+1)l} K_l F_k, \\
G_k R_l &= R_l G_k,
\end{aligned} \tag{15}$$

where the additional generators $\{K_i, T_i, R_i, \quad i \in Z\}$ must satisfy the relations (11) as in the one parameter deformation case and reduce to the unity operator when the deformation parameters p, q reduce to their classical values ($p = q = 1$)

In concluding we report that we established that it is possible to define a non trivial Hopf super-algebra structure on the deformed super Virasoro algebra, if we extend its deformed enveloping super-algebra with a set of additional even invertible elements. A consistence realisation of the additional even invertible elements $\{K_i, T_i, R_i, \quad i \in Z\}$ will be given elsewhere [40]

References

- [1] V. G. Drinfeld, Quantum groups, Proc. Int. Cong. Math.(Berkeley, 1986), Vol 1 798.
- [2] M. Jimbo, Lett. Math. Phys. 11 (1986) 247.
- [3] E. Abe, Hopf algebras (Cambridge U. Press, Cambridge, 1977).
- [4] P. P. Kulich and E. K. Sklyanin, In” Integrable quantum field theories”, J. Hietarinta and Montonen(eds); lect. Notes Phys, 151, (1982) 61-119
- [5] P. P. Kulich and N. Yu. Reshetikhin, lett. Math. Phys, 18, (1989) 143-149
- [6] S. M. Khoroshkim and V. N. Tolstoy Commun. Math. Phys, 141,(1991)599
- [7] P. P. Kulich and N. Yu. Reshetikhin, lett. Math. phys, 18, (1991) 599-617
- [8] R.Floresanini, V.P. Spiridonov and L. Vinet, Commun, Math. Phys 137, (1991) 149-160
- [9] M. Chaichain, P. P. Kulich and J. Lukiersski, Phys. Lett. B, 262, N.1, (1991) 43-48
- [10] M. Chaichain and P. P. Kulich, Phys. Lett. B234, No.1/2, (1990)72-80.
- [11] L. Pittner and P. Uray, J. Math, 36, (1995)944
- [12] P. P. Kulich , Quantum super algebras $osp(2,1)$, Preprint RIMS-615, Kyoto,(1988)
- [13] V. G. Kac, Funct. Anal. Appl.9, 263(1975)
- [14] V. G. Kac, Commun. Math. Phys. 53 31, (197)
- [15] M. Scheurnert, The Theory of Lie superalgebras, lecture Notes in Mathematics vol. 716(Springer, Berlin, 1979)
- [16] A. A. Belavin, A. M. Polyakov and A. B. Zamolodchikov, Nucl. Phys. B241 (1984)333

- [17] P. Fayet and S. Ferrara Phys. Rep 32 C (1977) 250.
- [18] P. Evan Niewvernhuizen Phys. Rep 86 C (1981) 189
- [19] M. B. Green, J. H. Schwarz and E. Witten, superstring theory, Cambridge Univ. Press, 1987
- [20] J. Schwarz Phys. Rep. 89 (1982) 223
- [21] E. Ragoucy and P. Sorba ,Inter. Jour. of Moder. Phys A, Vol. 7, No. 13 (1992) 2883-2971
- [22] A. Neveu and J. Schwarz, Nucl. Phys. B31 (1971) 86
- [23] P. Ramond, Phys. Rev. D3 (1971) 2415
- [24] N. Aizawa, T. Kobayashi and H-T. Sato, Notes on Curtright-Zachos Deformations of $osp(1,2)$ and super Virasoro algebras, hep-th/9706176.
- [25] T. Kobayashi and T. Uematsu, Phys. Lett. B306 (1993); Z. Phys. C56 (1992) 193.
- [26] T. Kobayashi, Z. Phys. C60 (1993) 101.
- [27] H. Hayashi, Commun. Math. Phys 127 (1990) 129.
- [28] R. Floreanini, P. Spiridinov and L. Vinet, Phys. Lett B 242 (1990) 383.
- [29] M. Mansour, Inter. Jour. Theo. Phys. Vol. 36, No. 12 (1997).
- [30] L. C. Biedenharn, J. Phys. A 22 (1989) L873.
- [31] A. J. Macfarlane, J. Phys. A 22 (1989) 4581.
- [32] T. L. Curtright and C. K. Zachos, Phys. Lett B 243 (1990) 237.
- [33] M. Chaichian, D. Ellinas and Z. Popowicz, Phys. Lett B 248 (1990) 95.
- [34] N. Aizawa and H. Sato Phys. Lett. B 256 (1991) 185.
- [35] H. Sato, Phys. Lett. B 415 (1997) 170.
- [36] W. S. Chung , Ilnuovo Cimento vol 109 B N.6 (1994).

- [37] M. Mansour and E. H. Tahri, Mod. Phys. Lett A, V 14 n 12 (1999) and ICTP-preprint IC/98/125,
- [38] M. Mansour and M. EL Falaki , Acta Physica Slovaca Vol 49 N6, 915-919.
- [39] M. Mansour and E. H. Tahri, (University of Rabat)-L.P.T-preprint (1998), to be submitted.
- [40] M.Mansour Work in progress